arXiv:1608.05500v1 [math.RT] 19 Aug 2016

THE BOUNDED SPHERICAL FUNCTIONS ON THE CARTAN MOTION GROUP AND GENERALIZATIONS FOR THE EIGENSPACES OF THE LAPLACIAN ON $\mathbb{R}^n$

JINGZHE XU

Abstract. The bounded spherical functions are determined for a real Cartan Motion group which is a generalization for the case when the Cartan Motion group is complex written by Sigurdur Helgason[1]. Also, I will do a further step of the Laplacian on $\mathbb{R}^n$. I consider the case when K is transitive on the spheres about 0 in $\mathbb{R}^n$,$n > 1$

1. Introduction

Consider a symmetric space $X = G/K$ of noncompact type, G being a connected noncompact semisimple lie group with finite center and K a maximal compact subgroup. Let $g = k + p$ be the corresponding Cartan decomposition, p being the orthocomplement of k relative to the killing form of g. Let $a \subset p$ be a maximal abelian subspace. Let G_0 be the Cartan Motion group. This group is defined as the semidirect product of K and p with respect to the adjoint action of K on p. The $X_0 = G_0/K$ is naturally identified with the Euclidean space p. The element $g_0 = (k, Y)$ actions on p by

$g_0(Y') = Ad(k)Y' + Y \ k \in K, Y, Y' \in p,$

So the algebra $\mathbb{D}(X_0)$ of G_0-invariant differential operators on X_0 is identified with the algebra of $Ad(K)$-invariant constant coefficient differential operators on p. The corresponding spherical functions on X_0 are given by $\psi_\lambda(Y) = \int_k e^{i\lambda(Ad(k)Y)}dk \ \lambda \in a_c^*$

and $\psi_\lambda = \psi_\mu$ if and only if λ and μ are W-conjugate. See e.g.[2],IV§4. Again, the maximal ideal space of $L^\natural(G_0)$ is up to W-invariance identified with the set of λ in a_c^* for which ψ_λ is bounded. Since ρ is relative to the curvature of G/K it is natural to expect the bounded ψ_λ to come from replacing $c(\rho)$ by the origin, where $c(\rho)$ is for the semisimple case also proved by Sigurdur Helgason [3]. In the words, ψ_λ is would be expected to be bounded if and only if λ is real, that is $\lambda \in a^*$. In [1], Sigurdur Helgason proved when G is complex, the spherical function ψ_λ on G_0 is bounded if and only if λ is real, i.e. $\lambda \in a^*$ mainly by using two results proved by Harish-Chandra [4] and [5]. In this paper, I use a different way to prove

1991 *Mathematics Subject Classification.* 22E46, 22E47.

Key words and phrases. Cartan Motion Group,spherical functions, eigenspaces of the Laplacian on $\mathbb{R}^n$.

when G is real, the spherical function ψ_λ on G_0 is bounded also if and only if λ is real. In this way, we generalize Sigurdur Helgason's results.

In Sigurdur Helgason's another paper [6], he considered Eigenspaces of the Laplacian on $\mathbb{R}^n$. Let L denote the usual Laplacian on $\mathbb{R}^n$ and for each $\lambda \in \mathbb{C}$ let $\mathcal{E}_\lambda(\mathbb{R}^n)$ denote the eigenspace $\mathcal{E}_\lambda(\mathbb{R}^n) = \{f \in \mathcal{E}(\mathbb{R}^n) \mid Lf = -\lambda^2 f\}$ with the topology induced by that of $\mathcal{E}(\mathbb{R}^n)$. Let G denote the group of all isometries of $\mathbb{R}^n$, and K the group of rotations $O(n)$. Sigurdur Helgason mainly proved the natural action of G on $\mathcal{E}_\lambda(\mathbb{R}^n)$ is irreducible if and only if $\lambda \neq 0$. I will prove when $K \subset O(n)$ is transitive on the spheres about 0 in $\mathbb{R}^n$,$n > 1$ instead of $O(n)$ and $G = K \rtimes \mathbb{R}^n$, the same results holds. In this way, we do a further step of this kind of problem. Meanwhile, I will specific all the groups K which is transitive on the spheres about 0 in $\mathbb{R}^n$.

Finally, according to [7], we know when $K \subset O(n)$ is transitively on the spheres about 0, the specific form of the spherical functions on $K \rtimes \mathbb{R}^n/K \cong \mathbb{R}^n$. Then I will give a estimation for it when $r \to \infty$.

2. The main theorem

The notion of induced spherical function mirrors the notion of induced representation. Let $Q \subset G$ be a closed subgroup such that K is transitive on G/Q, i.e.$G = KQ$,i.e.$G = QK$,i.e. Q is transitive on G/K. Let $\zeta : Q \to \mathbb{C}$ be spherical for $(Q, Q \bigcap K)$. The induced spherical function is $[Ind_Q^G(\zeta)](g) = \int_K \widetilde{\zeta}(gk)d\mu_K(k)$ where $\widetilde{\zeta}(kq) = \zeta(q)\Delta_{G/Q}(q)^{-\frac{1}{2}}$

Here $\Delta_{G/Q} : Q \to \mathbb{R}^n$ is the quotient of modular functions, $\Delta_{G/Q}(q) = \Delta_G(q)/\Delta_Q(q) = \Delta_Q(q)^{-1}$

Theorem 2.1. *Let $\lambda \in a_c^*$. Then $\psi_\lambda(Y)$ is the induced spherical function $Ind_p^G(\varphi_\lambda)$, where $\varphi_\lambda(Y) = e^{i\lambda(Y)}$ for every $Y \in a \subset p$.*

Proof. Apply above formula to φ_λ with $Q = P$. Since $G = K \rtimes p$ and p are unimodular, it says that the induced spherical function is given by $Ind_p^G(\varphi_\lambda)(k, Y) = \int_K \varphi_\lambda(Ad(k_0)Y)dk_0 = \int_K e^{i\lambda(Ad(k_0)Y)}dk_0 = \int_K e^{i\lambda(Ad(k)Y)}dk = \psi_\lambda(Y) = \psi_\lambda(k, Y)$ for $Y \in p, k \in K$. Note that $(k_0, 0)(k, Y) = (k_0 k, Ad(k_0)Y)$. □

We apply the Mackey little group method to G relative to its normal subgroup p:If ψ is an irreducible unitary representation of G,then it can be constructed (up to unitary equivalence) as follows: If $\varphi_\lambda(Y) = e^{i\lambda(Y)}$, where $\lambda \in p^*$, $Y \in p$, let $K_{\varphi_\lambda} = \{k \in K \mid \varphi_\lambda(Ad(k)Y) = \varphi_\lambda(Y)\ \forall Y \in p\}$. K_{φ_λ} is a closed subgroup of K. Let $G_{\varphi_\lambda} = K_{\varphi_\lambda} \rtimes p$. Write $\widetilde{\varphi_\lambda}$ for the extension of φ_λ to G_{φ_λ} given by $\widetilde{\varphi_\lambda}((k, Y)) = \varphi_\lambda(Y)$. If γ is an irreducible unitary representation of K_{φ_λ}, let $\widetilde{\gamma}$ denote its extension of G_{φ_λ} given by $\widetilde{\gamma}((k, Y)) = \gamma(k)$. Denote $\psi_{\varphi_\lambda,\gamma} = Ind_{G_{\varphi_\lambda}}^G(\widetilde{\varphi_\lambda} \otimes \widetilde{\gamma})$, then there exist choices of φ_λ and γ such that $\psi = \psi_{\varphi_\lambda,\gamma}$.

Theorem 2.2. *In the notation above, $\psi_{\varphi_\lambda,\gamma}$ has a K-fixed vector is given(up to scalar multiple by $u((k, Y)) = e^{-i\lambda(Ad(k^{-1})Y)}$, if $\varphi_\lambda = e^{i\lambda(Y)}$.*

Proof. The representation space H_ψ of $\psi = \psi_{\varphi_\lambda,\gamma}$ consists of all L^2 functions $f : G \to H_\psi$ such that $f(g^{'}(k^{'}, x^{'})) = \gamma(k^{'})^{-1}\varphi_\lambda(x^{'})^{-1}f(g^{'})$ for $g^{'} \in G$, $x^{'} \in p$, $k^{'} \in K_{\varphi_\lambda}$, and ψ acts by $(\psi(g)f)(g^{'}) = f(g^{-1}g^{'})$.

Now suppose that $0 \neq f \neq\in H_\psi$ is fixed under $\psi(K)$. If $k^{'} \in K_{\varphi_\lambda}$, then $\gamma(k^{'})f(1) = f(1)$. If $f(1) = 0$, then $f(G_{\varphi_\lambda}) = 0$ and K-invariance says $f = 0$, contrary to the assumption. Thus $f(1) \neq 0$ and irreducibility of γ forces γ to be trivial.

Conversely, if γ is trivial, then $u((k, Y)) = e^{-i\lambda(Ad(k^{-1})Y)}$ is a nonzero K-fixed vector in H_ψ. And it is the only one, up to scalar multiple, because any two K-fixed vectors must be proportional. □

Lemma 2.3. *In the notation above, $Ind_{G_{\varphi_\lambda}}^{G}(\widetilde{\varphi_\lambda})$ is unitary equivalent to the subrepresentation of $Ind_{p}^{G}(\widetilde{\varphi_\lambda})$ generated by the K-fixed unit vector $u((k, Y)) = e^{-i\lambda(Ad(k^{-1})Y)}$, if $\varphi_\lambda = e^{i\lambda(Y)}$.*

Theorem 2.4. *Let φ be a $(K \rtimes p, K)$-spheirical function. Then φ is positive definite if and only if it is of the form ψ_λ for some $\lambda \in a^*$. Further, if $\lambda, \lambda^{'} \in a^*$, then $\psi_\lambda = \psi_{\lambda'}$ if and only if $\lambda^{'} \in Ad(k)\lambda$.*

Proof. Let $\lambda \in a^*$. $\sum_{i,j} e^{i\lambda(Ad(k)(-Y_j+Y_i))}\overline{c_j}c_i = \sum_{i,j}(e^{-i\lambda(Ad(k)Y_j)}\overline{c_j})e^{i\lambda(Ad(k)Y_i)}c_i = (\sum_i e^{i\lambda(Ad(k)Y_i)}c_i)\overline{(\sum_j e^{i\lambda(Ad(k)Y_j)}c_j)} \geqslant 0$.

Since ψ_λ is a limit of non-negative linear combinations of positive definite functions on $\mathbb{R}^n$, so it is positive definite.

Now let φ be a positive definite (G, K)-spherical function. Let Π_φ be the associated irreducible unitary representation, and H_φ the representation space, such that there is a K-fixed unit vector $u_\varphi \in H_\varphi$ and let $\varphi(g) =< u_\varphi, \prod_\varphi(g)u_\varphi >$ for all $g \in G$. Following the discussion of the Mackey little group method, and Theorem 2.2 , we have $\varphi_\lambda(Y)$ for some $\lambda \in a^*, Y \in p$, s.t. Π_φ is unitarily equivalent to $Ind_{G_{\varphi_\lambda}}^{G}(\widetilde{\varphi_\lambda})$. Making the identification one, K-fixed unit vector in H_φ is given by $u((k, Y)) = e^{-i\lambda(Ad(k^{-1})Y)}$. We have $\lambda \in a*$ s.t.$\varphi_\lambda = e^{i\lambda(Y)}$ and from above several Theorems and Lemma, we compute:

$\varphi(Y) =< u, \Pi_\varphi(Y)u > =\varphi(Y) =< u, Ind_{G_{\varphi_\lambda}}^{G}(\widetilde{\varphi_\lambda})(Y)u >$

$=\varphi(Y) =< u, Ind_{G_p}^{G}(\varphi_\lambda)(Y)u >$

$=Ind_{G_p}^{G}(\varphi_\lambda)(Y)$

$=\psi_\lambda(Y)$

$=\int_K e^{i\lambda(Ad(k)Y)}dk$

For the second, if $\lambda^{'} = Ad(k_0)\lambda$ for some $k_0 \in K$, we have:

$\psi_{\lambda'}(Y) = \int_K e^{i\lambda^{'}(Ad(k)Y)}dk = \int_K e^{iAd(k_0)\lambda(Ad(k)Y)}dk =\int_K e^{i\lambda(Ad(k_0^{-1})Ad(k)Y)}dk = \int_K e^{i\lambda(Ad(k)Y)}dk = \psi_\lambda(Y)$.

Conversely, suppose that $\lambda^{'}, \lambda \in a^*$ with $\psi_{\lambda'} = \psi_\lambda$. Then (up to unitary equivalence)$Ind_{G_{\varphi_\lambda}}^{G}(\widetilde{\varphi_\lambda})= Ind_{G_{\varphi_{\lambda'}}}^{G}(\widetilde{\varphi_{\lambda'}})$. That gives us direct integral decompositions

$$\int_K^{\oplus} \psi_{Ad(k)\lambda} dk = Ind_{G_{\varphi_\lambda}}^G (\widetilde{\varphi_\lambda}) \mid p = Ind_{G_{\varphi_{\lambda'}}}^G (\widetilde{\varphi_{\lambda'}}) \mid p = \int_K^{\oplus} \psi_{Ad(k)\lambda'} dk \quad \square$$

Theorem 2.5. *If N is an n-step group with $n \geq 3$, then there are no Gelfand pairs (K, N), where $K \in Aut(N)$.*

Theorem 2.6. *We first consider K-spherical functions associated to a Gelfand pair (K, N).*

Suppose ϕ is a bounded K-spherical function on N. Then there is a $\pi \in \hat{N}$ and a unit vector $\xi \in H_\pi$ such that

$$\phi(x) = \int_K < \pi(k.x)\xi, \xi > dk \tag{2.7}$$

for each $x \in N$

Proof. Let $\lambda_\phi : L_K^1(N) \to \mathbb{C}$ be given by integration against ϕ.

Since $L^1(N)$ is a symmetric Banach *-algebra,[8],there is a representation $\bar{\pi}$ of $L^1(N)$ and a one-dimensional subspace H_ϕ of $H_{\bar{\pi}}$ such that $(\bar{\pi} \mid_{L_K^1(N)}, H_\phi)$ is equivalent to $(\lambda_\phi, \mathbb{C})$. As λ_ϕ is irreducible, the extension $\bar{\pi}$ is also irreducible(cf.[9]). Using approximate identities at each point of N, one can show that $\bar{\pi}$ is the integrated version of some $\pi \in \hat{N}$, with $H_\pi = H_{\bar{\pi}}$.

Choose $\xi \in H_\phi$ with $\|\xi\| = 1$. Then for each $f \in L_K^1(N)$, $\pi(f)\xi = \lambda_\phi(f)\xi$, so that

$$\begin{aligned} &< \phi, f >= \lambda_\phi(f) =< \pi(f)\xi, \xi > \\ &= \int_N f(x) < \pi(x)\xi, \xi > dx \\ &= \int_K \int_N f(k^{-1}.x) < \pi(x)\xi, \xi > dxdk \end{aligned} \tag{2.8}$$

since f is K-invariant

$$= \int_K \int_N f(k.x) < \pi(x)\xi, \xi > dxdk \tag{2.9}$$

Since ϕ is K-invariant, we change the order of integration and obtain

$$\phi(x) = \int_K < \pi(x)\xi, \xi > dk \tag{2.10}$$

$\square$

A complex-valued continous function ϕ on a locally compact group G is called positive definite if $\sum_{i,j=1}^n \phi(x_i^{-1}x_j)\alpha_i\overline{\alpha_j} \geq 0$ for all finite sets $x_1, \ldots, x_n$ of elements in G and any complex numbers $\alpha_1, \ldots, \alpha_n$.

Theorem 2.11. *For Gelfand pair (K, N), where N is at most 2-step nilpotent lie group, if ϕ is a bounded K-spherical function on N is and only if ϕ is positive definite.*

Proof. If ϕ is a bounded K-spherical function on N, for all finite sets $x_1, \ldots, x_n$ of elements in G and any complex numbers $\alpha_1, \ldots, \alpha_n$, we have:

$\sum_{i,j=1}^{n} \phi(x_i^{-1}x_j) = \int_K < \pi(k(x_i^{-1}x_j)\xi, \xi > dk = \int_K < \pi(k(x_i)^{-1}k(x_j)\xi, \xi > dk = \int_K < \pi(k(x_j)\xi, k(x_i)\xi > dk.$

Therefore, $\sum_{i,j=1}^{n} \phi(x_i^{-1}x_j)\alpha_i\overline{\alpha_j} = \sum_{i,j=1}^{n} \int_K < \overline{\alpha_j}\pi(k(x_j))\xi, \overline{\alpha_i}\pi(k(x_i))\xi > dk = \int_K < \sum_j \overline{\alpha_j}\pi(k(x_j))\xi, \sum_i \overline{\alpha_i}\pi(k(x_i))\xi > dk \geq 0.$

Conversely, if ϕ is a bounded K-spherical function on N then ϕ is positive definite. Let φ be a positive definite (G, K)-spherical function. Let Π_φ be the associated irreducible unitary representation, and H_φ the representation space, such that there is a K-fixed unit vector $u_\varphi \in H_\varphi$ and let $\varphi(g) = < u_\varphi, \prod_\varphi(g)u_\varphi >$ for all $g \in G$. Then we have $|\varphi(g)| = \left| < u_\varphi, \prod_\varphi(g)u_\varphi > \right|$

$\leq |< u_\varphi, u_\varphi >|^{\frac{1}{2}} \times \left| < \prod_\varphi(g)u_\varphi, \prod_\varphi(g)u_\varphi > \right|^{\frac{1}{2}} \leq |< u_\varphi, u_\varphi >| = 1$

Therefore, φ is bounded. □

Theorem 2.12. *In the notation just above, assume the group G real . The spherical function ψ_λ on G_0 is bounded if and only if λ is real,i.e.$\lambda \in a^*$.*

Proof. According to Theorem 2.4, we obtain The spherical function ψ_λ on G_0 is positive definite if and only if λ is real,i.e.$\lambda \in a^*$. According to Theorem 2.11, since p is abelian, we know that ψ_λ is a bounded K-spherical function on p is and only if ψ_λ is positive definite. Therefore, ψ_λ on G_0 is bounded if and only if λ is real,i.e.$\lambda \in a^*$. □

3. Generalizations for the Eigenspaces of the Laplacian on $\mathbb{R}^n$

Lemma 3.1. *[7] Let K be any closed subgroup of $O(n)$, if K is transitive on the spheres about 0 in $\mathbb{R}^n$, then $\mathcal{D}(G/K) = \mathbb{C}[\triangle]$, algebra of polynomials in the Laplace-Beltrami operator $\triangle = -\sum \partial^2/\partial x_i^2$.*

Proof. It is clear that $\mathbb{C}[\triangle] \subset \mathcal{D}(G/K)$. Now let $D \in \mathcal{D}(G/K)$ be of order m. Then the mth order symbol of D is a pollynomial of pure degree m constant on spheres about 0 in $\mathbb{R}^n$, in other words a multiple cr^m with m even and $r^2 = \sum x_i^2$. Now $D - (c(-\triangle)^{m/2}) \in \mathcal{D}(G/K)$ and $D - (c(-\triangle)^{m/2})$ has order $< m$. By induction on the order, $D - (c(-\triangle)^{m/2}) \in \mathcal{D}(G/K)$, so we have $D \in \mathcal{D}(G/K)$. □

Let L denote the usual Laplacian on $\mathbb{R}^n$ and for each $\lambda \in \mathbb{C}$ let $\mathcal{E}_\lambda(\mathbb{R}^n)$ denote the eigenspace

$$\mathcal{E}_\lambda(\mathbb{R}^n) = \{f \in \mathcal{E}(\mathbb{R}^n) \mid Lf = -\lambda^2 f\} \tag{3.2}$$

with the topogy induced by that of $\mathcal{E}(\mathbb{R}^n)$. Let $G = K \rtimes \mathbb{R}^n$ and K is the closed subgroup of $O(n)$ as well as acting transitive on the spheres about 0 in $\mathbb{R}^n$.

Theorem 3.3. *The natural action of G on $\mathcal{E}_\lambda(\mathbb{R}^n)$ is irreducible if and only if $\lambda \neq 0$.*

Proof. It is clear that each function

$$f(x) = \int_{S^{n-1}} e^{i\lambda(x,w)} F(w)dw,\ F \in L^2(S^{n-1}), \tag{3.4}$$

lies in $\mathcal{E}_\lambda(\mathbb{R}^n)$; here $(,)$ denotes the usual inner product on $\mathbb{R}^n$ and dw the normalized volume element. □

Lemma 3.5. *Let $\lambda \neq 0$. Then the mapping $F \to f$ defined by (3.4) is one-to-one.*

Proof. Let $p(\zeta) = p(\zeta_1, \cdots \zeta_n)$ be a polynomial and D the corresponding constant coefficient differential operator on $\mathbb{R}^n$ such that

$$\int_{\mathbb{R}^n} e^{i(x,\zeta)} D_x(e^{-(\frac{1}{2})|x|^2}) = p(\zeta)e^{-(\frac{1}{2})(\zeta_1^2+\cdots+\zeta_n^2)} \tag{3.6}$$

for$\zeta \in \mathbb{C}^n$. If $f \equiv= 0$ in (1) we deduce from (3.6) that

$$\int_{S^{n-1}} p(\lambda w_1, \cdots, \lambda w_n)F(w)dw = 0 \tag{3.7}$$

Since $\lambda \neq 0$, this implies $F \equiv 0$. □

Lemma 3.8. *Let $\lambda \neq 0$. The K-finite solutions f of the equation $Lf = -\lambda^2 f$ are precisely*

$$f(x) = \int_{S^{n-1}} e^{i\lambda(x,w)} F(w)dw \tag{3.9}$$

where F is a K-finite function on S^{n-1}.

Proof. Let δ be an irreducible representation of K and if Σ is any sphere in $\mathbb{R}^n$ with center at 0 let $\mathcal{E}_\delta(\Sigma)$ denote the space of K-finite functions in $\mathcal{E}(\Sigma)$ of type δ. We know from Lemma 1.5 p.134 in [10] that if Σ is suitably chosen each function $f \mid \Sigma$ toΣ. With F and f as in (3.4) it follows that the maps

$F \to f \mid \Sigma,\ F \to f\ F \in L^2(S^{n-1})$

are one-to-one and commute with the action of K. For reasons of dimensionality, the first must therefore map $\mathcal{E}_\delta(S^{n-1})$ onto $\mathcal{E}_\delta(\Sigma)$. The lemma now follows. □

For $\lambda \neq 0$ let $\mathcal{H}_\lambda$ denote the space of functions f as defined in (3.4); $\mathcal{H}_\lambda$ is a Hilbert space if the norm of f is the L^2 norm of F on S^{n-1}.

Lemma 3.10. *Let $\lambda \neq 0$. Then the space $\mathcal{H}_\lambda$ is dense in $\mathcal{E}_\lambda(\mathbb{R}^n)$.*

Proof. Each eigenfunction of L can be expanded in a convergent series of K-finite eigenfunctions (cf. Sect. 5 [6]) so the lemma follows from Lemma 3.8. □

We can now prove Theorem 3.3. We first prove that G acts irreducibly on $\mathcal{H}_\lambda$. Let $V \neq 0$ be a closed invariant subspace of $\mathcal{H}_\lambda$. Then there exists an $h \in V$ such that $h(0) = 1$. We write

$$h(x) = \int_{S^{n-1}} e^{i\lambda(x,w)} H(w) dw \tag{3.11}$$

and the average $h^\natural(x) = \int_K h(k.x)dk$ is then

$$h^\natural(x) = \varphi_\lambda(x) = \int_{S^{n-1}} e^{i\lambda(x,w)} dw. \tag{3.12}$$

If f in (3.4) lies in the annihilator V^0 of V the functions F and H are orthogonal on S^{n-1}. Since V^0 is K-invariant this remains true for H replaced by its integral over K, in other words φ_λ belongs to the double annihilator $(V^0)^0 = V$. Now, since V is invariant under translations it follows that for each $t \in \mathbb{R}$ the function

$$x \to \int_{S^{n-1}} e^{i\lambda(x,w)} e^{i\lambda(t,w)} dw \tag{3.13}$$

belongs to V. But then Lemma 3.5 shows that the annihilator of V in $\mathcal{H}_\lambda$ is 0, whence the irreducibility of G on $\mathcal{H}_\lambda$.

Passing now to $\mathcal{E}_\lambda$ let $V \subset \mathcal{E}_\lambda$ be a closed invariant subspace. Then $V \cap \mathcal{H}_\lambda$ is an invariant subspace of $\mathcal{H}_\lambda$; Schwartz' inequality shows easily that it is closed. Thus, by the above, $V \subset \mathcal{E}_\lambda$ is $\{0\}$ or $\mathcal{H}_\lambda$. In the second case $V = \mathcal{E}_\lambda$ by Lemma 3.10. In the first case consider for each $f \in V$ the convergent expansion

$$f = \sum_{\delta \in \hat{K}} \alpha_\delta * f \tag{3.14}$$

where $\alpha_\delta = d(\delta)\chi_\delta^*$ and

$$(\alpha_\delta * f)(x) = \int_K \alpha_\delta(k) f(k^{-1}.x) dk \tag{3.15}$$

χ_δ being the character of δ.Then $\alpha_\delta * f \in \mathcal{H}_\lambda$ by Lemma 3.8. Let $V^0 \subset \mathcal{E}'(\mathbb{R}^n)$ be the annihilator of V. Then V^0 is G-invariant and if $T \in V^0$,

$$\int_{\mathbb{R}^n} (\alpha_\delta * f)(x) dT(x) = \int_K \alpha_\delta(k) \int_{\mathbb{R}^n} f(x) dT(k.x) dk \tag{3.16}$$

so $\alpha_\delta * f$ belongs to the double annihilator $(V^0)^0 = V$. Thus $\alpha_\delta * f \in V \cap \mathcal{H}_\lambda = \{0\}$ so, by (3.6),$f = 0$. Thus $V = \{0\}$ so the proof is finished.

4. A estimation for some spherical functions and the groups K

From [7], we know if K is transitive on the spheres about 0 in $\mathbb{R}^n$, then the spherical function on $K \rtimes \mathbb{R}^n \simeq \mathbb{R}^n$ is of the form: $\varphi_s(r) = \varphi(r,s) = \int_{S^{n-1}} e^{s(\xi,x)} d\sigma(\xi) = \frac{\Gamma(\frac{n}{2})}{\sqrt{\pi}\Gamma\frac{n-1}{2}} \int_0^\pi e^{sr\cos\theta} \sin^{n-2}\theta d\theta$.

Where s is a complex number, and $r = \|x\| = \sqrt{x_1^2 + \ldots + x_n^2}$, S^{n-1} is the unit sphere in $\mathbb{R}^n$ and σ the normalized surface measure on S^{n-1}.

If Res=0 then it follows from above equation that $|\varphi_s(x)| \leq 1$ for all $x \in G$.

Clearly,$\varphi_s = \varphi_{-s}$. We just need to consider the case Re$s \geq 0$.

Theorem 4.1. *If Re$s > 0$, we have* $\varphi(r,s) \sim \frac{\Gamma(\frac{n}{2})2^{\frac{n-3}{2}}}{\sqrt{\pi}} \frac{e^{sr}}{(sr)^{\frac{n-1}{2}}}$, *when* $r \to \infty$.

Proof. From above equation, we know that, by an elementary substitution,
$\varphi_s(r) = \varphi(r,s) = \frac{\Gamma(\frac{n}{2})}{\sqrt{\pi}\Gamma(\frac{n-1}{2})} \int_{-1}^{1} e^{srt}(1-t^2)^{\frac{n-3}{2}} dt$
and, setting $t = 1 - \frac{u}{r}$, we obtain
$\varphi(r,s) = \frac{\Gamma(\frac{n}{2})}{\sqrt{\pi}(\frac{n-1}{2})} \frac{e^{sr}}{r^{\frac{n-1}{2}}} \int_0^{2r} e^{-su} u^{\frac{n-3}{2}} (2 - \frac{u}{r})^{\frac{n-3}{2}} du$
For Re$s > 0$, we get, using Lebesgue's dominated convergence theorem,
$\lim_{r\to\infty} \int_0^{2r} e^{-su} u^{\frac{n-3}{2}} (2 - \frac{u}{r})^{\frac{n-3}{2}} du = \frac{\Gamma(\frac{n-1}{2})2^{\frac{n-3}{2}}}{s^{\frac{n-1}{2}}}$ □

Finally, I will give all the possible K, which is transitive on the spheres about 0 on $\mathbb{R}^n$, $n > 1$. [11]

When K is transitive on the spheres about 0 in $\mathbb{R}^n$,$n > 1$, its identity compoment K^0 is also transitive, and $K = K^0F$ where F is a finite subgroup of the normalizer $N_{O(n)}(K^0)$. The possibilities for K^0 are as follows:

(1)$n > 1$ and $K^0 = SO(n)$,
(2)$n = 2m$ and (i)$K^0 = SU(m)$ or (ii)$U(m)$,
(3)$n = 4m$ and (i)$K^0 = Sp(m)$ or (ii)$Sp(m).U(1)$ or $Sp(m).Sp(1)$,
(4)$n = 7$ and K^0 is the exceptional group G_2,
(5)$n = 8$ and $K^0 = Spin(7)$, and
(6)$n = 16$ and $K^0 = Spin(9)$.

In case (1), $N_{O(n)}(K^0) = O(n)$, so the relevant choices for F are $\{I\}$ and $\{I, -I\}$, so K is either $SO(n)$ or $O(n)$.

In case (2)(i),$N_{O(n)}(K^0) = U(m) \bigcup \alpha U(m)$ where α is complex conjugation of $\mathbb{C}^m$ over $\mathbb{R}^m$. The relevant choices for F are the finite subgroups of $U(1) \bigcup \alpha U(1)$ where $U(1)$ consists of the unitary scalar matrices $e^{ix}I$, x real. Those are the cyclic groups $\mathbb{Z}_l = \{e^{2\pi ik/l}I\}$ of order $l \geq 1$ and the dihedral groups $\mathbb{D}_l = \mathbb{Z}_l \bigcup \alpha\mathbb{Z}_l$, so K is a group $SU(m)\mathbb{Z}_l$ or $SU(m)\mathbb{D}_l$. In the case (2)(ii) the relevant possibilities for F are $\{I\}$ and $\{\alpha, I\}$, so K is either $U(m)$ or $U(m) \bigcup \alpha U(m)$.

In case (3)i,(3)(iii),(4),(5),(6), K^0 has no outer automorphism, so we may take F in the centralizer $Z_{O(n)}(K^0)$. Thus in the case (3)(i), F can be any subgroup of $Sp(1)$, in the other words, a cyclic group $\mathbb{Z}_l$ of order l, a binary dihedral group $\mathbb{D}_l^*$ of order $4l$, a binary tetrahedral group $\mathbb{T}^*$ of order 24, a binary octahedral group $\mathbb{O}^*$ of order 48, or a binary icosahedral group $\mathbb{I}^*$ of order 60. Thus K is a group $Sp(m)\mathbb{Z}_l$,$Sp(m)\mathbb{D}_l^*$,$Sp(m)\mathbb{T}^*$, $Sp(m)\mathbb{O}^*$ or $Sp(m)\mathbb{I}^*$. In case (3)(ii) the relevant possibilities for F are $\{I\}$ and $\{\beta, I\}$, where the $U(1)$ factor of K^0 consists of all quaternion scalar multiplications by complex numbers e^{ix}, x is real, as in the case (2), and

β is quaternion scalar multiplication by j.Thus K is either $Sp(m)U(1)$ or $(Sp(m)U(1))\bigcup(Sp(m)U(1))\beta$. In case (3)(iii), K^0 is its own $O(n)$-centralizer so $F = \{I\}$ and $K = Sp(m)Sp(1)$.

In case (4),(5),(6),K^0 is absolutely irreducible on $\mathbb{R}^n$, so relevant F would have to consist of real scalars. As G_2 does not contain $-I$ we see that the relevant F for case (4) are $\{I\}$ and $\{I, -I\}$, resulting in $K = G_2$ and $K = G_2\bigcup(-I)G_2$. Both $Spin(7)$ and $Spin(9)$ do contain $-I$, so F is trivial in case (5),(6).That gives $K = Spin(7)$ in case (5) and $K = Spin(9)$ for case (6).

References

[1] Helgason, Sigurdur *The Bounded Spherical Functions on the Cartan motion group*,Arxiv ID: 1503.07598.

[2] S. Helgason, *Goups and Geometric Analysis:Integral Geometry,Invariant Differential Operators,and Spherical Functions*, Academic Press. (1984).

[3] S. Helgason and K. Johnson, The bounded spherical functions on symmetric spaces Adv. Math. 3 (1969), 586593.

[4] Harish-Chandra, *Differential operators on a semisimple Lie algebra*, Amer. J. Math. 79 (1957), 241310

[5] Harish-Chandra *Spherical functions on a semisimple Lie group I*, Amer. J. Math 80 (1958), 241310.

[6] Helgason, Sigurdur, *Eigenspaces of the Laplacian; integral representations and irreducibility*, Journal of Functional Analysis, 1974, Vol.17(3), pp.328-353

[7] Wolf, Joseph A, *Spherical functions on Euclidean space.*, J. Funct. Anal. 239 (2006) No.1, 127-136.

[8] H. Leptin *On group algebras of nilpotent groups*, Studia Math. 47 (1973), 37-49

[9] M. Naimark, *Normed rings*, Wolters-Noordhoff, 1970.

[10] Helgason, Sigurdur, *A duality for symmetric spaces with applications to group representations,*, Advan. Math. 5 (1970), l-154.

(Xu) Department of Mathematics, Hong Kong University of Science and technology, Clear Water Bay, Kowloon, Hong Kong SAR, China

E-mail address: jxuad@ust.hk